

Classification of regular maps of negative prime Euler characteristic

Antonio Breda d'Azevedo

Departamento de Matematica
Universidade de Aveiro
Aveiro, Portugal

Roman Nedela

Institute of Mathematics
Slovak Academy of Science
Banská Bystrica, Slovakia

Jozef Širáň

Department of Mathematics, SvF
Slovak University of Technology
Bratislava, Slovakia

Abstract

We give a classification of all regular maps on nonorientable surfaces with a negative odd prime Euler characteristic (equivalently, on nonorientable surfaces of genus $p + 2$ where p is an odd prime). A consequence of our classification is that there are no regular maps on nonorientable surfaces of genus $p + 2$ where p is a prime such that $p \equiv 1 \pmod{12}$ and $p \neq 13$.

1 Introduction

A regular map is an embedding of a graph in a compact, connected surface, such that the automorphism group of the embedding acts regularly on flags (edges with a longitudinal and a transverse direction). If the surface is orientable and one requires only the orientation-preserving automorphism group of the embedding to be regular on arcs (edges with longitudinal direction) then the map is orientably regular.

The ubiquitous five platonic solids give the most well known examples of regular maps. Non-spherical regular maps appeared in form of stellated polyhedra in the work of Kepler [21] as early as in 1619. Origins of more current interest in the study of regular maps go back to the late 19th century and were brought up by two independent streams of research. In connection with Heawood's work on map colourings that eventually led to the rise of modern topological graph theory, Heffter [15] constructed orientably regular embeddings of complete graphs of prime order. On the other hand, certain three-valent regular maps on a surface of genus three were studied by Klein [22] and Dyck [10] in connection with constructions of automorphic functions on surfaces. In the beginning of the 20th century, regular maps appeared as a form of a geometric representation of groups in Burnside's monograph [5]. The first systematic treatment of regular maps was due to Brahana [3] and was later developed on the levels of both geometry and combinatorial group theory by Coxeter and Moser [8]. Modern foundations of the theory of maps on orientable surfaces can be found in Jones and Singerman [19]. An analogous theory for maps on nonorientable surfaces was outlined by Bryant and Singerman [4].

Links between the theory of regular and orientably regular maps, group theory, hyperbolic geometry, and complex functions have been known for more than a century. In the orientable case they can be briefly summed up as follows. Automorphism groups of orientably regular maps on surfaces of genus at least two are quotients of triangle groups by torsion-free normal subgroups of finite index. The same quotient construction can be used to endow maps with complex structure and hyperbolic geometry from the Poincaré complex upper half-plane on which the triangle groups act. An additional boost to the study of regular maps was lately given by a part of Grothendieck's programme [14] to investigate the absolute Galois group by means of its action on maps. For more reading about these exciting connections we recommend the survey papers by Jones [18] and by Jones and Singerman [20].

In the light of the above facts, classification of regular and orientably regular maps is one of the important problems whose solution would be in the interest of (and may find applications in) disciplines such as topological graph theory, group theory, hyperbolic geometry, and the theory of Riemann surfaces. The problem has been approached in three natural ways: classification by underlying graphs, by automorphism groups, and by supporting

surfaces. Since our focus is on classification by surfaces we mention only three most significant results in the first two approaches. An abstract characterization of graphs underlying regular and orientably regular maps was given by Gardiner, Nedela, Širáň and Škovič [12]. A classification of orientably regular embeddings of complete graphs can be found in James and Jones [17]. For maps with given group, Sah [25] gave a classification of all orientably regular maps with automorphism group isomorphic to $PSL(2, q)$.

We now briefly survey regular maps on a fixed surface. The only surfaces supporting infinitely many regular maps are a sphere, a projective plane, and a torus. The infinitude in the first two cases is due to trivial maps whose underlying graphs are cycles, dipoles, and semistars. In contrast, there is no regular map on a Klein bottle. For the remaining cases it follows from the Hurwitz bound (see Tucker [27]) that the order of the automorphism group of a regular map on a surface of negative Euler characteristic χ cannot exceed -84χ . This implies that the number of regular maps on such surfaces is automatically finite. A strengthening of the above bound to $-8\chi + 16$ for infinitely many χ follows from the work of Accola [1] or MacLachlan [23].

Regular and orientably regular maps on orientable surfaces of genus at most seven together with regular maps on nonorientable surfaces of genus at most eight have been classified by the late 1980's. This was an outcome of effort of a multitude of authors over a considerable time span (see Brähana [3], Coxeter and Moser [8], Sherk [26], Garbe [11], Bergau and Garbe [2] and references therein). For genus two or more the main method was relation-chasing, supported by certain combinatorial arguments. With the help of the low index subgroup algorithm applied to certain finitely presented groups related to triangle groups, Conder and Dobcsányi [6] recently gave a computer assisted classification of all regular and orientably regular maps on orientable surfaces of genus at most 15 and a classification of all regular maps on nonorientable surfaces of genus at most 30. Thus, complete lists of regular and orientably regular maps were available only for a finite number of surfaces at the time of submission of this article.

A particularly interesting open question was whether there exist infinitely many nonorientable surfaces supporting no regular map at all. Conder and Everitt [7] constructed a variety of infinite families of regular maps on nonorientable surfaces, covering about 75 per cent of characteristics. On the other hand, Wilson and Breda [29] proved that among all nonorientable surfaces of genus at most 52 only those of genus 2, 3, 18, 24, 27, 39 and 48 do not

support a regular map.

This paper represents a breakthrough in the nonorientable regular map classification problem. We derive a complete classification of all nonorientable regular maps with negative odd prime Euler characteristic – or, equivalently, regular maps on nonorientable surfaces of genus $p + 2$ where p is an odd prime. As a by-product we also obtain an affirmative answer to the above question. To be able to state our main result in a condensed form, for $p \equiv -1 \pmod{4}$ we denote by $\nu(p)$ the number of pairs of positive integers (j, l) such that $j > l \geq 3$, both j and l are odd, and $(j - 1)(l - 1) = p + 1$.

Theorem 1 *Let p be an odd prime, $p \neq 7, 13$, and let \mathcal{N}_{p+2} be a nonorientable surface of Euler characteristic $-p$ (and hence of genus $p + 2$).*

- (1) *If $p \equiv 1 \pmod{12}$ then there is no regular map on \mathcal{N}_{p+2} .*
- (2) *If $p \equiv 5 \pmod{12}$ then, up to isomorphism and duality, there is exactly one regular map on \mathcal{N}_{p+2} .*
- (3) *If $p \equiv -5 \pmod{12}$ then, up to isomorphism and duality, there are $\nu(p)$ regular maps on \mathcal{N}_{p+2} .*
- (4) *If $p \equiv -1 \pmod{12}$ then, up to isomorphism and duality, \mathcal{N}_{p+2} supports exactly $\nu(p) + 1$ regular maps.*

A more detailed statement of this result in group-theoretic language will appear later as Theorem 2 in Section 2, preceded by essentials of the theory of regular maps. In Section 3 we reduce the problem of classifying regular maps on nonorientable surfaces of Euler characteristic $-p$ (p a prime) by proving that the orders of the corresponding groups are severely limited if $p \geq 29$. Using a powerful result of Gorenstein and Walter [13] that characterizes groups with dihedral Sylow 2-subgroups, in Sections 4 and 5 we further restrict the class of candidates for automorphism groups of our regular maps. The final Section 6 contains a proof of Theorem 2 and concluding remarks.

2 Regular maps and groups

A precise definition of a map requires introducing flags or their equivalents such as oriented forms [28], or edges with longitudinal and transverse direction or blades [4]. This in general requires discussing a number of degenerate cases. The object of our interest, however, are maps that are regular and on

nonorientable surfaces. For such maps, *flags* can be identified with (topological) triangles whose three distinguished points are a vertex, the “center” of an edge incident with the vertex, and the “center” of a face incident with both the vertex and the edge. Each face bounded by an m -gon is thus subdivided into $2m$ flags.

It is well known that the automorphism group of a regular map can be generated by three involutions x, y, z reflecting a fixed flag in its three sides and sending it to the three incident flags. One may choose the notation in such a way that the compositions $yz, zx,$ and xy are rotations of the map about the vertex, the center of the face, and the center of the edge associated with the fixed flag. If k and m are the vertex valence and the face length of M we say that M is of *type* $\{m, k\}$. In such a case the automorphism group of M has a presentation of the form $Aut(M) = \langle x, y, z \mid x^2 = y^2 = z^2 = (yz)^k = (zx)^m = (xy)^2 = \dots = 1 \rangle$ where dots indicate a possible presence of additional independent relators. We emphasize that in all group presentations in this article, exponents will be assumed to be *true orders* of the corresponding elements.

Motivated by the above, we define a $(k, m, 2)$ -*group* to be any finite group $G = \langle x, y, z \rangle$ generated by an ordered triple (x, y, z) of involutions and presented in the form

$$G = \langle x, y, z \mid x^2 = y^2 = z^2 = (yz)^k = (zx)^m = (xy)^2 = \dots = 1 \rangle .$$

It follows that $(k, m, 2)$ -groups are finite torsion-free quotients of the extended $(k, m, 2)$ -triangle group [4]. We will say that two $(k, m, 2)$ -groups $G = \langle x, y, z \rangle$ and $G' = \langle x', y', z' \rangle$ are *congruent* if there is a group isomorphism from G onto G' taking x to x' , y to y' , and z to z' .

Any $(k, m, 2)$ -group $G = \langle x, y, z \rangle$ acts on a (unique) closed surface as the automorphism group of a regular map $M = (G; x, y, z)$ of type $\{m, k\}$. Flags of M are elements of G , and edges, vertices and faces of M are right cosets of the dihedral subgroups $\langle x, y \rangle$, $\langle y, z \rangle$, and $\langle z, x \rangle$ of G , respectively. Mutual incidence of the map elements is given by nonempty intersection, and G acts on M as map automorphism group by left multiplication. This, in principle, enables one to identify regular maps of type $\{m, k\}$ with $(k, m, 2)$ -groups; a precise statement will be given in Proposition 1.

Let $G = \langle x, y, z \rangle$ and $G' = \langle x', y', z' \rangle$ be a pair of $(k, m, 2)$ -groups and let $M = (G; x, y, z)$ and $M' = (G'; x', y', z')$ be the corresponding regular maps. Then, M and M' are *isomorphic* if the groups G and G' are congruent. Further, the maps M and M' are *dual* of each other if there is a group

isomorphism from G onto G' which sends x onto y' , y onto x' , and z onto z' . It is easy to see that these definitions exactly correspond to the traditional concepts of map isomorphism and map duality.

Let \mathcal{S} be the supporting surface of a regular map $M = (G; x, y, z)$ and let $\chi(\mathcal{S})$ be the Euler characteristic of \mathcal{S} . The Euler characteristics $\chi(G)$ of the $(k, m, 2)$ -group $G = \langle x, y, z \rangle$ and $\chi(M)$ of the map M are defined by $\chi(G) = \chi(M) = \chi(\mathcal{S})$. They can be obtained from Euler's formula by substituting $|G|/2k$, $|G|/4$, and $|G|/2m$ for the number of vertices, edges, and faces, respectively, giving $\chi(M) = \chi(G) = (1/k + 1/m - 1/2)|G|/2$.

We have the following obvious but important consequence of the above considerations.

Proposition 1 *Regular maps on a surface of Euler characteristic $\chi < 2$ are, up to isomorphism and duality, in a one-to-one correspondence with congruence classes of $(k, m, 2)$ -groups, $k \geq m$, of Euler characteristic χ .*

Given a $(k, m, 2)$ -group $G = \langle x, y, z \rangle$, we will be using throughout the notation $r = yz$ and $s = zx$. The supporting surface \mathcal{S} of the regular map $M = (G; x, y, z)$ is nonorientable if and only if $G = \langle r, s \rangle$. Note that if G has an odd Euler characteristic (which is going to be the case in this article) then \mathcal{S} is automatically nonorientable. Therefore we will also work with the *reduced* presentation $G = \langle r, s \mid r^k = s^m = (rs)^2 = \dots = 1 \rangle$. In general one should be aware of a loss of information when “forgetting” about the original presentation of G in terms of x, y, z . The reason is that there may be several non-equivalent ways to choose the involutions x, y, z such that $r = yz$ and $s = zx$, leading to non-congruent groups and hence non-isomorphic regular maps whose automorphism groups have the same reduced presentation. But our proofs will show that this will never be the case with the maps and groups considered here.

We will use the standard notation Z_n , D_{2n} , $PGL(2, q)$ and $PSL(2, q)$ for a cyclic group of order n , dihedral group of order $2n$, and 2-dimensional projective general and special linear group over $GF(q)$, respectively. In addition we need to introduce two special classes of groups. For any odd prime p such that $p \equiv -1 \pmod{3}$ let G_p be the $(p+4, 4, 2)$ -group of order $8(p+4)$ with reduced presentation

$$G_p = \langle r, s \mid r^{p+4} = s^4 = (rs)^2 = sr^3s^{-1}r^3 = 1 \rangle .$$

Before defining our second special class of groups, recall that for any prime p such that $p \equiv -1 \pmod{4}$ we denoted by $\nu(p)$ the number of pairs (j, l) of

odd integers j and l such that $j > l \geq 3$ and $(j-1)(l-1) = p+1$. Now, for any pair (j, l) as above let $G_{j,l}$ be the $(2j, 2l, 2)$ -group of order $4jl$ with reduced presentation

$$G_{j,l} = \langle r, s \mid r^{2j} = s^{2l} = (rs)^2 = rs^2r^{-1}s^2 = sr^2s^{-1}r^2 = 1 \rangle .$$

The $\nu(p)$ groups $G_{j,l}$ obtained this way will be referred to as groups *associated* with p . Note that for $p \equiv -1 \pmod{4}$ and $p \geq 7$ we always have $\nu(p) \geq 1$ because of the pair $j = (p+3)/2$ and $l = 3$. We will show later in Section 5 that the groups G_p and $G_{j,l}$ are well defined. In particular, G_p and $G_{j,l}$ are abstractly isomorphic to a product $Z_{p+4} \cdot D_8$ and to a direct product $D_{2j} \times D_{2l}$, respectively.

We are now ready to present the extended version of our main result, stated in terms of $(k, m, 2)$ -groups and their reduced presentations. The reason for using group-theoretic language is twofold. Firstly, all our proofs are purely group-theoretic. Secondly, the translation back to regular maps is straightforward by Proposition 1.

Theorem 2 *Let p be an odd prime and let $n(p)$ be the number of pairwise non-congruent $(k, m, 2)$ -groups $G = \langle x, y, z \rangle = \langle r, s \rangle$ of Euler characteristic $-p$ such that $k \geq m$.*

- (1) *If $p \equiv 1 \pmod{12}$ and $p \neq 13$, then $n(p) = 0$.*
- (2) *We have $n(13) = 1$, and the corresponding group is a $(7, 3, 2)$ -group of order 1092 isomorphic to $PSL(2, 13)$ with reduced presentation*

$$\langle r, s \mid r^7 = s^3 = (rs)^2 = ((sr^{-2})^4 sr^3)^2 = 1 \rangle .$$
- (3) *If $p \equiv 5 \pmod{12}$, then $n(p) = 1$. The unique representative here is the $(p+4, 4, 2)$ -group G_p .*
- (4) *If $p \equiv 7 \pmod{12}$ and $p \neq 7$, then $n(p) = \nu(p)$ and the corresponding groups are the $(2j, 2l, 2)$ -groups $G_{j,l}$ associated with p .*
- (5) *We have $n(7) = 3$, where one of the groups is the $(10, 6, 2)$ -group $G_{5,3}$ and the other two groups are the $(8, 3, 2)$ -groups of order 336 isomorphic to $PGL(2, 7)$ with reduced presentations*

$$\langle r, s \mid r^8 = s^3 = (rs)^2 = (sr^{-2})^4 = 1 \rangle \quad \text{and}$$

$$\langle r, s \mid r^8 = s^3 = (rs)^2 = ((sr^{-2})^2 r^{-2})^2 = 1 \rangle .$$

- (6) *If $p \equiv -1 \pmod{12}$, then $n(p) = \nu(p) + 1$, the groups being the $\nu(p)$ groups $G_{j,l}$ associated with p , together with the group G_p .*

For completeness we mention that for $k \geq m$ there are exactly eight non-congruent $(k, m, 2)$ -groups of Euler characteristic -2 , six of which are automorphism groups of regular maps on orientable surfaces [6].

3 Preliminary reduction results

Let G be a $(k, m, 2)$ -group of Euler characteristic $-p$ where p is an odd prime. Since $p = -\chi(G) = (1/2 - 1/k - 1/m)|G|/2$, it follows that $1/k + 1/m < 1/2$; the $(k, m, 2)$ -groups satisfying this inequality will be called *hyperbolic*. Using $\mu(k, m) = 4km/(km - 2k - 2m)$ we see that the order of a hyperbolic $(k, m, 2)$ -group G of Euler characteristic $-p$ is given by $|G| = \mu(k, m)p$.

It is known that for positive integers k and m such that $1/k + 1/m < 1/2$ we have $\mu(k, m) \leq 84$, with equality if and only if $\{k, m\} = \{3, 7\}$. The next three largest values of $\mu(k, m)$ are 48 for $\{k, m\} = \{3, 8\}$, 40 for $\{k, m\} = \{4, 5\}$, and 36 for $\{k, m\} = \{3, 9\}$. We note that the reciprocal of $\mu(k, m)$ for $1/k + 1/m < 1/2$ is equal to the area of the fundamental region for the action of the extended $(k, m, 2)$ -triangle group as a group of isometries in a hyperbolic plane.

Since a computer assisted characterization of all $(k, m, 2)$ -groups of (not necessarily prime) Euler characteristic χ is known [6] for all $\chi \geq -28$, it is sufficient to assume that $p \geq 29$. We use this remark to derive a divisibility condition for the denominator of $\mu(k, m)$.

Proposition 2 *Let G be a hyperbolic $(k, m, 2)$ -group of Euler characteristic $-p$ where p is a prime. If $p \geq 29$ then $(km - 2k - 2m) = pt$ for a positive integer t dividing $4km$, and $|G| = 4km/t$.*

Proof. Assume the contrary and let a prime number $p \geq 29$ be not a divisor of $km - 2k - 2m$. Since $|G| = \mu(k, m)p$ we see that $\mu(k, m)$ must be an integer, and as G contains the Klein four-group it follows that $4|\mu(k, m)$. We first claim that p^2 is not a divisor of $|G|$. Indeed, in the opposite case p would have to divide $\mu(k, m)$ and so $p \leq \mu(k, m)/4 \leq 21$, a contradiction. In particular, p divides neither k nor m , and the Sylow p -subgroups of G are isomorphic to Z_p .

Suppose now that $H \simeq Z_p$ is a unique Sylow p -subgroup of G . Then H is normal in G , and since neither k nor m have p as a factor it follows that the quotient group G/H is a $(k, m, 2)$ -group again. From $\mu(k, m) = |G|/p =$

$|G/H| = -\chi(G/H)\mu(k, m)$ we see that $\chi(G/H) = -1$. It is known, however, that there are no $(k, m, 2)$ -groups of Euler characteristic -1 (see e.g. [6]). Consequently, G contains more than one Sylow p -subgroup.

By Sylow theorems, the number of Sylow p -subgroups in G is equal to $np + 1$ for some $n \geq 1$ such that $np + 1$ is a divisor of $|G| = \mu(k, m)p$. This readily shows that $p^2 < |G|$, or $p < \mu(k, m)$, which implies that $29 \leq p \leq 83$. But it is easily checked that there are exactly four sets $\{k, m\}$ with $4|\mu(k, m)$ that satisfy the inequalities $29 \leq p < \mu(k, m)$, namely, the four sets listed before the statement of Proposition 2.

Out of these sets we can quickly eliminate $\{k, m\} = \{4, 5\}$ with $\mu(4, 5) = 40$ and $\{k, m\} = \{3, 9\}$ with $\mu(3, 9) = 36$, since $np + 1$ divides neither $40p$ nor $36p$ for any prime $p \geq 29$ and any $n \geq 1$. The only two possibilities for $\{k, m\} = \{3, 7\}$ with $\mu(3, 7) = 84$ are $p = 83$ and $p = 41$, giving a potential of two $(2, 3, 7)$ -groups of orders $84 \cdot 83$ and $84 \cdot 41$. But both can be excluded since by Sylow theorems they contain a unique (and hence normal) cyclic Sylow 7-subgroup H , collapsing in G/H the generator of G of order 7 and thus producing a quotient of too a small order.

It remains to eliminate the case when $\{k, m\} = \{3, 8\}$, with $\mu(3, 8) = 48$. We may without loss of generality assume that $G = \langle r, s \mid r^3 = s^8 = (rs)^2 = \dots = 1 \rangle$. There must be exactly 48 Sylow 47-subgroups in G , occupying a total of 47^2 elements. This shows that elements of order 2^i , $1 \leq i \leq 4$, must be between the remaining 47 elements. Consider the (dihedral) Sylow 2-subgroup $H = \langle s, z \rangle \simeq D_{16}$ of order 16 in G . If $H \neq g^{-1}Hg$ for some $g \in G$ of order 47, then all the 47 conjugates of H by powers of g would be pairwise distinct, giving too many elements of order a power of 2 in G . It follows that all the 47^2 elements of order 47 normalize H , and hence H is normal in G . But then the quotient is easily seen to collapse.

It follows that $km - 2k - 2m = pt$ for a positive integer t . From $|G| = \mu(k, m)p$ we then obtain $|G| = 4km/t$ and so t must be a divisor of $4km$. \square

For any given prime $p \geq 29$, Proposition 2 greatly restricts orders and parameters k, m of the possible $(k, m, 2)$ -groups G of Euler characteristic $-p$: One has to have $|G| = 4km/t$ for some t such that $km - 2k - 2m = tp$. In the remaining part of this section we show that $t \in \{1, 2, 4\}$.

Lemma 1 *Let $G = \langle x, y, z \rangle = \langle r, s \rangle$ be a $(k, m, 2)$ -group with $k, m \geq 3$ and let $L = \langle y, z \rangle \cap \langle z, x \rangle$. Then, L is isomorphic to $Z_2 \times Z_2$ if and only if both k and m are even and $r^{k/2}s^{m/2} \in \langle z \rangle$; in all other cases $L \simeq Z_2$.*

Proof. Since $z \in L$ the group L contains a subgroup isomorphic to Z_2 . Assume that $\langle z \rangle$ is a proper subgroup of L . Being a subgroup of both $\langle r, z \rangle$ and $\langle s, z \rangle$ the group L must be dihedral. Let j be the smallest positive divisor of k such that $L = \langle r^j, z \rangle$. If r^j is not an involution then it must be in the cyclic subgroup of $\langle s, z \rangle$, that is, $r^j = s^l$ for some l . But from $G = \langle r, s \rangle$ we see that the element $b = r^j = s^l$ is central in G ; in particular, $yby = b$. On the other hand, conjugation by y inverts $b = r^j = (yz)^j$, and so $b = b^{-1}$, contrary to the fact that r^j was assumed to be non-involutory. Since $k, m \geq 3$, we have $r^\alpha \notin \{y, z\}$ for $1 \leq \alpha \leq k-1$ and $s^\beta \notin \{x, z\}$ for $1 \leq \beta \leq m-1$. It follows that k must be even and $L = \langle r^{k/2}, z \rangle$; by the symmetry of the argument we also have m even and $L = \langle s^{m/2}, z \rangle$. The lemma follows. \square

Lemma 2 *Let G and L be as in Lemma 1. If $km - 2k - 2m = tp$, then $t \leq |L|$. In particular, $t \in \{1, 2, 4\}$. Moreover:*

- (1) $t = 4$ if and only if $|L| = 4$, and in this case $G = \langle r \rangle \langle s \rangle$ if and only if $z = r^{k/2} s^{m/2}$;
- (2) $t = 2$ if and only if exactly one of k, m is even, with $(k, m) = 1$, and then $G = \langle r \rangle \langle z, x \rangle = \langle y, z \rangle \langle s \rangle$;
- (3) $t = 1$ if and only if both k and m are odd and $(k, m) = 1$.

Proof. The assumption $km - 2k - 2m = tp$ implies that $|G| = 4km/t$. By the formula $|AB| \cdot |A \cap B| = |A| \cdot |B|$, valid for any two subgroups A, B of G , we have $4km/t = |G| \geq |\langle y, z \rangle \langle z, x \rangle| = |\langle y, z \rangle| |\langle z, x \rangle| / |L| = 4km/|L|$. This shows that $t \leq |L|$; in particular, by Lemma 1 we have $t \leq 4$, with $t = 4$ implying that $|L| = 4$. Conversely, if $|L| = 4$ then, invoking Lemma 1 again, k and m must be even, say, $k = 2j$ and $m = 2l$. From $t \leq 4$ and $tp = km - 2k - 2m = 4jl - 4j - 4l$ we see that $t = 4$. This shows that $t < 4$ if and only if $|L| = 2$, and therefore $t \neq 3$. Also, if $t = |L| = 4$ we have $km = |G| \geq |\langle r \rangle \langle s \rangle| = km/|L_o|$ where $L_o = \langle r \rangle \cap \langle s \rangle$. From the proof of Lemma 1 it is obvious that $|L_o| \leq 2$, with L_o trivial if and only if $z = r^{k/2} s^{m/2}$. This proves part (1). Parts (2) and (3) follow immediately from the equation $km - 2k - 2m = tp$ with help of calculations similar to the presented ones. \square

For any integer n let π_n , the *parity* of n , be 1 or 2 according as n is odd or even. We note that in terms of parity the three parameters k, m and t from Lemma 2 are related by $t = \pi_k \pi_m$.

4 Hyperbolic $(k, m, 2)$ -groups with dihedral Sylow two-subgroups

Before we consider individual cases for $t \in \{1, 2, 4\}$ we prove a statement that further substantially restricts potential candidates for our groups. Let $G = \langle x, y, z \rangle = \langle r, s \rangle$ be a hyperbolic $(k, m, 2)$ -group, where $k, m \geq 3$ are orders of r and s , respectively. Throughout this section we will assume that k is odd.

Let $|G| = 4km/\pi_m$ where π_m is the parity of m introduced earlier. Let c be the largest integer such that 2^c divides $|G|$. If $\pi_m = 1$, then $c = 2$ and $\langle x, y \rangle \simeq Z_2 \times Z_2$ is a Sylow 2-subgroup of G . If $\pi_m = 2$, then 2^c divides $2m$, and since k is odd we see that G has a dihedral Sylow 2-subgroup contained in $\langle z, x \rangle$. It follows that in any case G has a dihedral Sylow 2-subgroup.

In what follows we will use the the following deep result of Gorenstein and Walter (see [13], Part I, Page 85, Theorem 1): If G is a group with a dihedral Sylow 2-subgroup and if $O(G)$ is the (unique) maximal normal subgroup of G of odd order, then $G/O(G)$ is isomorphic to either

- (a) a Sylow 2-subgroup of G , or
- (b) the alternating group A_7 , or
- (c) a subgroup of $Aut(PSL(2, q))$ containing $PSL(2, q)$, q odd.

It is well known that $Aut(PSL(2, q))$, q an n -th power of an odd prime, is a semidirect product of $PGL(2, q)$ with a cyclic group of order n that comes from the Galois action on the standard matrix representation of $PGL(2, q)$. It follows that any subgroup of $Aut(PSL(2, q))$ that contains $PSL(2, q)$ has the form of a split extension $H:Z_b$ where H is either $PSL(2, q)$ or $PGL(2, q)$ and b is a divisor of n . Let $\delta = 1/2$ or $\delta = 1$ according as $H \simeq PSL(2, q)$ or $H \simeq PGL(2, q)$, and let ω_1 and ω_2 be the largest order and the second largest order of an element in $H:Z_b$. Combining the above facts with Dickson's classification [9] of possible orders of elements of $PSL(2, q)$ and $PGL(2, q)$ we conclude the following: If $n \geq 2$ then $\omega_1 = \delta b(q + 1)$ and $\omega_2 = \delta b(q - 1)$, and if $n = 1$ then $\omega_1, \omega_2 \in \{\delta(q + 1), q\}$.

We are now in position to prove a structural result about $(k, m, 2)$ -groups with dihedral Sylow 2-subgroups.

Proposition 3 *Let $G = \langle x, y, z \rangle = \langle r, s \rangle$ be a $(k, m, 2)$ -group and let $|G| = 4km/\pi_m$ where $k, m \geq 3$, $(k, m) = 1$, and k is odd. Then $G/O(G)$ is isomorphic to S_4 or A_5 according as m is even or odd.*

Proof. As we saw, G has a dihedral Sylow 2-subgroup. By the result of Gorenstein and Walter [13], one of (a), (b), and (c) above holds for $G/O(G)$. We first exclude (a) and (b).

In the case (a) the subgroup $O(G)$ is a normal 2-complement in G . It follows that G is a semidirect product of $O(G)$ and a Sylow subgroup S_2 of G . In particular, the element r of odd order k must be in $O(G)$. Let $s = gh$ with $g \in O(G)$ and $h \in S_2$ and let d be the order of h . Note that $d < |S_2|$ since the Sylow 2-subgroup is dihedral. But then, $|\langle r, s \rangle| \leq |O(G)| \cdot d < |O(G)||S_2| = |G|$, a contradiction. The case (b) is easily excluded by observing that the order of A_7 is much larger than $4km/\pi_m = |G|$ where k and m are orders of two elements of A_7 . Therefore, for $G/O(G)$ we have the situation described in (c), and it remains to show that $G/O(G)$ is isomorphic to $PGL(2, 3) \simeq S_4$ or to $PSL(2, 5) \simeq A_5$ depending on the parity of m .

Keeping to the notation introduced before the statement of Proposition 3, we have concluded from (c) that $G/O(G) \simeq H:Z_b$ where H is either $PSL(2, q)$ or $PGL(2, q)$, q and n -th power of an odd prime, and $b|n$. Denoting the cosets of a subgroup of G by bars above the corresponding elements, let k' and m' be the orders of \bar{r} and $\bar{s} = \bar{z}\bar{x}$ in $G/O(G) = \langle \bar{r} \rangle \langle \bar{z}, \bar{x} \rangle$; clearly $k' \neq m'$. Now, if $n \geq 2$ then $\delta q(q^2 - 1)b = |H:Z_b| = |G/O(G)| \leq 4k'm'/\pi_m \leq 4\delta b(q+1)\delta b(q-1)$. This implies that $q \leq 4\delta b \leq 4n$, which is absurd for any n -th power q of any odd prime if $n \geq 2$. For $n = 1$ we obtain $\delta q(q^2 - 1) = |H| = |G/O(G)| \leq 4k'm'/\pi_m \leq 4\delta q(q+1)/\pi_m$. If m is even, this gives a contradiction for all odd $q \geq 5$ but is feasible if $q = 3$ (for both values of δ); if m is odd then we have a contradiction for all odd $q \geq 7$ but feasibility for $q = 3$ or $q = 5$.

For even m it follows that $G/O(G)$ is isomorphic to either $PSL(2, 3) \simeq A_4$ or $PGL(2, 3) \simeq S_4$. The group A_4 , however, cannot be generated by three involutions and hence $G/O(G) \simeq S_4$, as claimed. For odd m and $q = 3$ we only have the groups $PSL(2, 3) \simeq A_4$ and $PGL(2, 3) \simeq S_4$ as candidates for $G/O(G)$, neither of which can be generated by two elements of coprime odd orders. Finally, if m is odd and $q = 5$ then $G/O(G)$ can only be isomorphic to $PSL(2, 5) \simeq A_5$ or to $PGL(2, 5) \simeq S_5$. But the only two non-trivial coprime odd orders here are 3 and 5 and the corresponding permutations in S_5 are even, showing that $G/O(G) \simeq A_5$ in this case. \square

5 Further reduction results

We begin with analyzing the individual cases $t = 1, 2,$ and 4 . If $t = 1$ then, by part (3) of Lemma 2 we have $|G| = 4km$ with both k and m odd, and $(k, m) = 1$. In this situation, Proposition 3 can be strengthened as follows.

Proposition 4 *Let $G = \langle x, y, z \rangle = \langle r, s \rangle$ be a $(k, m, 2)$ -group with $|G| = 4km$ where $k, m \geq 3$, both k and m are odd, and $(k, m) = 1$. Then $G \simeq A_5$.*

Proof. Let G be a counterexample of the smallest order. From Proposition 3 it follows that $G/O(G) \simeq A_5$, and so $|O(G)| \geq 3$. We first show that no non-trivial proper subgroup of $O(G)$ is normal in G . Indeed, if K was such a subgroup then G/K would satisfy the assumptions of the theorem, and from $60 < |G/K| < |G|$ we would obtain a contradiction with minimality of $|G|$.

Without loss of generality we may suppose that $G/O(G) = \langle \bar{r}, \bar{s} \mid \bar{r}^5 = \bar{s}^3 = (\bar{r}\bar{s})^2 = 1 \rangle \simeq A_5$, so that $k = 5k'$ and $m = 3m'$ for some odd and coprime k', m' such that $\max\{k', m'\} \geq 3$. Since $|G| = 4km = 60k'm'$, we see that $|O(G)| = k'm'$. Further, observe that $r^5, s^3 \in O(G)$, and since the subgroups $\langle r^5 \rangle$ and $\langle s^3 \rangle$ intersect just trivially, it follows that $O(G) = \langle r^5 \rangle \langle s^3 \rangle$. Then, by Ito's theorem [16], the commutator subgroup $K = [O(G), O(G)]$ is a proper characteristic subgroup of $O(G)$, and so K is normal in G . By the observation in the preceding paragraph we conclude that $K = 1$, that is, $O(G)$ is abelian; since $|O(G)| = k'm'$ is a product of two coprime numbers, $O(G)$ is cyclic. Now, if an odd prime p' is a proper divisor of $k'm'$, then $O(G)$ contains a unique (and hence characteristic) subgroup of order p' that would be normal in G , contradiction. Therefore $O(G)$ is cyclic of odd prime order p' , and either $k = 5p'$ and $m = 3$, or $k = 5$ and $m = 3p'$.

To finish the proof we just deal with the case $k = 5p'$ and $m = 3$ as the argument is analogous for the other case. Now, $O(G) = \langle r^5 \rangle$, and since $O(G)$ is normal in G we have $xr^5x = r^{-5i}$ for some i ; since x is an involution we have $i^2 \equiv 1 \pmod{p'}$. But then, $sr^5s^{-1} = r^{5i}$, and this remains valid when conjugating r^5 by any odd power of s . From $G = \langle r, s \rangle$ and from the order of s being odd it follows that x can be written as a word $w = w(r, s)$ over the alphabet $\{r, s\}$ in which the letter s appears an odd number of times. Due to this fact we obtain $r^{-5i} = xr^5x = wr^5w^{-1} = r^{5i}$, a contradiction with $r^{5p'} = 1$ and $i^2 \equiv 1 \pmod{p'}$. \square

Let us now consider the situation when $t = 2$. By part (2) of Lemma 2 we know that $G = \langle r \rangle \langle z, x \rangle = \langle y, z \rangle \langle s \rangle$. From the same source it follows that exactly one of k and m is even, and $(k, m) = 1$. Without loss of generality we may assume that m is even.

Proposition 5 *Let $G = \langle x, y, z \rangle = \langle r, s \rangle$ be a hyperbolic $(k, m, 2)$ -group such that $|G| = 2km$ where k is odd, m is even, and $(k, m) = 1$. Then, $k = 3j$ with j odd, $m = 4$, and $G = \langle r \rangle \langle z, x \rangle \simeq Z_k \cdot D_8$, with reduced presentation*

$$\langle r, s \mid r^k = s^4 = (rs)^2 = sr^3s^{-1}r^3 = 1 \rangle .$$

Proof. By Proposition 3 we know that $G/O(G) \simeq S_4$. Since $G = \langle r \rangle \langle z, x \rangle$, the group $G/O(G)$ must also be a product of a cyclic group of odd order and a dihedral (or a cyclic) group. It is easy to see that S_4 can be this way decomposed in an essentially unique way as $S_4 = \langle \bar{r} \rangle \langle \bar{z}, \bar{x} \rangle \simeq Z_3 \cdot D_8$. The relations for the dihedral group D_8 transform to $\bar{z}^2 = \bar{x}^2 = (\bar{z}\bar{x})^4$. This means that z^2 , x^2 and $(zx)^4$ are all in $O(G)$. Since $O(G)$ has odd order, this is possible only if $\langle z, x \rangle \simeq D_8$. In particular, $m = 4$, and from $|G| = 2km = 8k$ we see that $G \simeq Z_k \cdot D_8$.

It remains to address the presentation of G . From $\bar{r}^3 = 1$ we have $r^3 \in O(G)$, and so $k = 3j$ for some odd j . We now have $|G| = 2km = 24j = |S_4||O(G)|$, which shows that here $O(G)$ is a cyclic group of order j generated by r^3 . By normality of $O(G)$ in G we have $xr^3x = r^{3i}$ (and hence $sr^3s^{-1} = r^{-3i}$ as well) for some i such that $i^2 \equiv 1 \pmod{j}$. Let $x = w(r, s)$ where $w = w(r, s)$ is a word in r and s . If s occurs in w an odd number of times, then $r^{3i} = xr^3x = wr^3w = r^{-3i}$ since $i^2 \equiv 1 \pmod{j}$, which (as in the preceding proof) quickly contradicts the fact that r has odd order. Therefore, s occurs in w an even number of times, and by the same token we have $r^{3i} = xr^3x = wr^3w = r^3$, implying that $i = 1$. \square

Finally, we are ready to deal with the case $t = 4$.

Proposition 6 *Let $G = \langle x, y, z \rangle = \langle r, s \rangle$ be a hyperbolic $(k, m, 2)$ -group with $|G| = km$ where $(k, m) \leq 2$. Then k and m are even, $k/2$ and $m/2$ are odd, and $G = \langle y, r^2 \rangle \times \langle x, s^2 \rangle \simeq D_k \times D_m$ has reduced presentation*

$$\langle r, s \mid r^{2j} = s^{2l} = (rs)^2 = rs^2r^{-1}s^2 = sr^2s^{-1}r^2 = 1 \rangle .$$

Proof. Invoking part (1) of Lemma 2 we see that $|G| = km$ is equivalent to $|L| = t = 4$, with $G = \langle y, z \rangle \langle z, x \rangle$. First, assume that $G = \langle r \rangle \langle s \rangle$, which (by Lemma 2 again) is equivalent to $k = 2j$, $m = 2l$, and $z = r^j s^l$.

By Ito's theorem [16], the commutator subgroup $G' = [G, G]$ of G is abelian. Note that both $r^2 = [y, z]$ and $s^2 = [z, x]$ are in G' . Clearly $(j, l) = 1$; we show that both j and l are odd. Indeed, if, say, j is even, then l must be odd. Right multiplication of both sides of $z = r^j s^l$ by s then gives $x = r^j s^{l+1} \in G'$. In particular, as both x and s^2 belong to the *abelian* G' , we have $x s^2 x = s^2$. But conjugation by x obviously inverts s^2 , and so $s^2 = s^{-2}$ and hence $l \leq 2$, contradiction.

Having both j, l odd we see that $rzs = r^{j+1} s^{l+1} \in G'$. Since $rxs = yzx = rxs$ is in the *abelian* subgroup G' , we have $(ys)s^2 = s^2(ys)$ and $(rx)r^2 = r^2(rx)$. It follows that y commutes with s^2 and x with r^2 . Since conjugation by y and x inverts r^2 and s^2 , respectively, we see that G is a direct product of the dihedral subgroups $\langle y, r^2 \rangle \simeq D_k$ and $\langle x, s^2 \rangle \simeq D_m$. The last two relators in the presentation of G are equivalent with y and x commuting with s^2 and r^2 , respectively, and the completeness of the presentation is easy to check.

To finish the proof we have to analyze the situation when $|G| = km$ and $G \neq \langle r \rangle \langle s \rangle$. From Lemma 1 and part (1) of Lemma 2 it is obvious that this is the case if and only if $k = 2j$, $m = 2l$, and $b = r^j = s^l$ is a central involution of G . Then, the quotient group $G/\langle b \rangle$ of order $km/2 = 2jl$ where $(j, l) = 1$, has presentation $\bar{r}^j = \bar{s}^l = (\bar{r}\bar{s})^2 = \dots = 1$. Observe that $j, l \geq 3$; otherwise one of j, l would be equal to 2 and then the three relations in the preceding presentation would define a dihedral group of order jl , contradiction. By part (2) of Lemma 2 applied to the $(j, l, 2)$ -group $G/\langle b \rangle$ of order $2jl$ we see that exactly one of j, l is even; without loss of generality let j be odd. From Proposition 5 we then see that $j = 3i$ for some odd i , $l = 4$, and $G/\langle b \rangle = \langle \bar{r}, \bar{s} \mid \bar{r}^{3i} = \bar{s}^4 = (\bar{r}\bar{s})^2 = \bar{s}\bar{r}^3 s^{-1} \bar{r}^3 = 1 \rangle \simeq Z_j \cdot D_8$. Moreover, centrality of b in G and presentation of $G/\langle b \rangle$ allows us to conclude that the original group G must have had a *full* presentation of the form $r^{6i} = s^8 = (rs)^2 = r^{3i} s^4 = xr^3 x r^{-3} r^{3\alpha} = 1$ where $\alpha = 0$ or $\alpha = i$. This shows that, in any case, $\langle r^6 \rangle$ is a normal subgroup of G of (odd) order i . The corresponding quotient group $G/\langle r^6 \rangle$ of order $|G|/i = 48$ then must be a $(6, 8, 2)$ -group of Euler characteristic $\chi(G/\langle r^6 \rangle) = (1/6 + 1/8 - 1/2) \cdot 24 = -5$. But by the list of Conder and Dobcsányi [6], no such group exists. \square

6 Proof of the main result and remarks

With the help of results established in the previous section we are in position to prove our main result.

Proof of Theorem 2. Let $G = \langle x, y, z \rangle = \langle r, s \rangle$ be a $(k, m, 2)$ -group of characteristic p where p is an odd prime and $k \geq m$. We know that such a group must be hyperbolic. The part of the list of groups in the statement of Theorem 2 corresponding to $p \leq 23$ is extracted from the classification of $(k, m, 2)$ -groups of Euler characteristic at least -28 , given in [6]. We may therefore assume that $p \geq 29$. By Proposition 2 and Lemma 2 we have $km - 2k - 2m = tp$ and $|G| = 4km/t$ where $t \in \{1, 2, 4\}$.

If $t = 1$ then we have $km - 2k - 2m = p$ and hence both k and m are odd and $(k, m) = 1$. By Proposition 4, up to isomorphism there is a unique such $(k, m, 2)$ -group of order $4km$, namely, the $(5, 3, 2)$ -group isomorphic to A_5 . But this group has Euler characteristic 1, a contradiction.

Now let $t = 2$. From $km - 2k - 2m = 2p$ it follows that exactly one of k, m is odd and $(k, m) = 1$. Proposition 5 then shows that (up to isomorphism) we have a unique $(k, m, 2)$ -group G of order $2km$ here, with $k = 3j$ for an odd j , and $m = 4$. Invoking the equation $km - 2k - 2m = 2p$ again we see that this is possible only if $p \equiv -1 \pmod{3}$ and $k = p + 4$. Therefore, G is the $(p + 4, 4, 2)$ -group G_p that appears in parts (3) and (6) of Theorem 2.

In the last case when $t = 4$, from $km - 2k - 2m = 4p$ we obtain $(k, m) \leq 2$. According to Proposition 6 we have $k = 2j$ and $m = 2l$ where $j > l$, both j and l are odd, and $(j, l) = 1$. Then, $km - 2k - 2m = 4p$ is equivalent with $(j - 1)(l - 1) = p + 1$ and hence $p \equiv -1 \pmod{4}$. Further, by Proposition 6, for each such pair (j, l) there is (up to isomorphism) a unique $(2j, 2l, 2)$ -group of order $km = 4jl$, namely, the group $G_{j,l}$. Recalling the function ν introduced in Section 2 we have a total of $\nu(p)$ non-congruent groups in this case, featuring in parts (4) and (6) of Theorem 2.

Thus, for $p \geq 29$, $(k, m, 2)$ -groups of a negative prime Euler characteristic $-p$ exist if and only if $p \equiv -1 \pmod{3}$ or $p \equiv -1 \pmod{4}$. It follows that for primes $p \geq 29$ there are no $(k, m, 2)$ -groups of Euler characteristic $-p$ if $p \equiv 1 \pmod{12}$, which is part (1) of Theorem 2. Parts (3), (4) and (6) of Theorem 2 for $p \geq 29$ are immediate consequences of the foregoing analysis.

Uniqueness of the $(k, m, 2)$ -groups whose *reduced* presentations are given as in Theorem 2 follows from our proof in Sections 4 and 5. An independent argument can be given by observing that the centers of our groups are trivial

and quoting Theorem 7.7 from [24] that can be restated as follows: If a group $G = \langle r, s \mid r^k = s^m = (rs)^2 = \dots = 1 \rangle$ has center of an odd order, then G has (up to congruence) a unique representation as a $(k, m, 2)$ -group. \square

We conclude with a few remarks regarding the map-theoretic interpretation of our main result.

From the presentation of the group G_p for prime $p \equiv -1 \pmod{3}$ it is clear that G_p contains a cyclic normal subgroup of order $(p+4)/3$ generated by r^3 . The quotient $G_p/\langle r^3 \rangle \simeq \langle \bar{r}, \bar{s} \mid \bar{r}^3 = \bar{s}^4 = (\bar{r}\bar{s})^2 = 1 \rangle$ is the automorphism group of a regular embedding M of a complete graph of order four in a projective plane. It follows that the regular maps corresponding to the groups G_p are cyclic normal $(p+4)/3$ -fold covers of M , with branch points of order $(p+4)/3$ at each of the four vertices of M . An analogous statement can be made about the dual maps. We note that the groups G_p appear also in [23] in connection with bounds for the number of automorphisms of a compact Riemann surface.

Similarly, for primes $p \equiv -1 \pmod{4}$ the groups $G_{j,l}$ associated with p contain a normal subgroup isomorphic to $Z_j \times Z_l$ generated by r^2 and s^2 . The reduced presentation $\langle \bar{r}, \bar{s} \mid \bar{r}^2 = \bar{s}^2 = (\bar{r}\bar{s})^2 = 1 \rangle \simeq Z_2 \times Z_2$ of the resulting quotient group represents a regular projective-planar embedding M' of a loop with a single vertex. We conclude that the $\nu(p)$ regular maps corresponding to the groups $G_{j,l}$ are normal $Z_j \times Z_l$ -covers of M' , with the two branch points being the vertex (of order j) and the center of the single face (of order l). More statements of this type can be generated by considering other normal subgroups of $G_{j,l}$.

Regular maps corresponding to the groups of Theorem 2 can be lifted from nonorientable supporting surfaces \mathcal{N}_{p+2} to their orientable double covers. Our reduced presentations thus automatically give a classification of *antipodal* regular maps on orientable surfaces of genus $p+1$ for prime p . This may be beneficial for possible future projects regarding classification of regular maps on orientable surfaces. More details about antipodality can be found in Nedela and Škoviera [24].

Finally, we note that by [29] there are no regular maps on nonorientable surfaces of Euler characteristic -16 , -22 , -25 and -46 , which are values not covered by part (1) of Theorem 2. The problem of determining all nonorientable surfaces that do not support any regular map remains open.

Acknowledgement. The authors thank the Department of Mathematics of the University of Aveiro and the Research Unit “Matemática e Aplicações” for supporting this project. Research of the second and the third authors was partially supported by the Slovak Grant Agency VEGA; the third author was also sponsored by the U.S.-Slovak Science and Technology Joint Fund under Project Number 020/2001.

References

- [1] R. D. M. Accola, On the number of automorphisms of a closed Riemann surfaces, *Trans. Amer. Math. Soc.* **131** (1968), 398–408.
- [2] P. Bergau and D. Garbe, Non-orientable and orientable regular maps, in: Proceedings of “Groups-Korea 1988”, *Lect. Notes Math.* 1398, Springer (1989), 29–42.
- [3] H. R. Brahana, Regular maps and their groups, *Amer. J. Math.* **49** (1927), 268–284.
- [4] R. P. Bryant, D. Singerman, Foundations of the theory of maps on surfaces with boundary. *Quart. J. Math. Oxford Ser. (2)* **36** (1985), no. 141, 17–41.
- [5] W. Burnside, “Theory of Groups of Finite Order”, Cambridge Univ. Press, 1911.
- [6] M. Conder and P. Dobcsányi, Determination of all regular maps of small genus, *J. Combinat. Theory Ser. B* **81** (2001), 224–242.
- [7] M. Conder and B. Everitt, Regular maps on non-orientable surfaces, *Geom. Dedicata* **56** (1995), 209–219.
- [8] H. S. M. Coxeter and W. O. J. Moser, “Generators and Relations for Discrete Groups”, 4th Ed., Springer-Verlag, Berlin, 1984.
- [9] E. Dickson, “Linear groups with an exposition of Galois field theory”, 1901; Dover Publ., 1958.

- [10] W. Dyck, Über Aufstellung und Untersuchung von Gruppe und Irrationalität regularer Riemannscher Flächen, *Math. Ann.* **17** (1880), 473–508.
- [11] D. Garbe, Über die regulären Zerlegungen geschlossener orientierbarer Flächen, *J. Reine Angew. Math.* **237** (1969), 39–55.
- [12] A. Gardiner, R. Nedela, J. Širáň and M. Škoviera, Characterization of graphs which underlie regular maps on closed surfaces, *J. London Math. Soc.* (2) **59** (1999) No. 1, 100–108.
- [13] D. Gorenstein and J. H. Walter, The characterization of finite groups with dihedral Sylow 2-subgroups, I, II, III, *J. of Algebra* **2** (1965), 85–151, 218–270, 334–393.
- [14] A. Grothendieck, “Esquisse d’un programme”, preprint, Montpellier (1984).
- [15] L. Heffter, Über metazyklische Gruppen und Nachbarconfigurationen, *Math. Ann.* **50** (1898) 261–268.
- [16] N. Ito, Über das Produkt von zwei abelschen Gruppen, *Math. Z.* **62** (1955), 400–401.
- [17] L. D. James and G. A. Jones, Regular orientable imbeddings of complete graphs, *J. Combinat. Theory Ser. B* **39** (1985), 353–367.
- [18] G. A. Jones, Maps on surfaces and Galois groups, *Math. Slovaca* **47** (1997), 1–33.
- [19] G. A. Jones and D. Singerman, Theory of maps on orientable surfaces, *Proc. London Math. Soc.* (3) **37** (1978), 273–307.
- [20] G. A. Jones and D. Singerman, Belyĭ functions, hypermaps, and Galois groups, *Bull. London Math. Soc.* **28** (1996), 561–590.
- [21] J. Kepler, “The harmony of the world” (translation from the Latin “*Harmonice Mundi*”, 1619), *Memoirs Amer. Philos. Soc.* **209**, American Philosophical Society, Philadelphia, PA, 1997.
- [22] F. Klein, Über die Transformation siebenter Ordnung der elliptischen Functionen, *Math. Ann.* **14** (1879), 428–471.

- [23] C. MacLachlan, A bound for the number of automorphisms of a compact Riemann surface, *J. London Math. Soc.* **44** (1969), 265–272.
- [24] R. Nedela and M. Škoviera, Exponents of orientable maps, *Proc. London Math. Soc.* (3) **75** (1997), 1–31.
- [25] C. H. Sah, Groups related to compact Riemann surfaces, *Acta Math.* **123** (1969), 13–42.
- [26] F. A. Sherk, The regular maps on a surface of genus three, *Canad. J. Math.* **11** (1959) 452–480.
- [27] T. W. Tucker, Finite groups acting on surfaces and the genus of a group, *J. Combinat. Theory Ser. B* **34** (1983) No. 1, 82–98.
- [28] W. T. Tutte, What is a map? in “New Directions in Graph Theory” (F. Harary, Ed.), Acad. Press, 1973, 309–325.
- [29] S. Wilson and A. Breda D’Azevedo, Surfaces with no regular hypermaps, to appear in *Discrete Math.*