

Regular t -balanced Cayley maps from Abelian groups

Martin Mačaj

GEMS 2009

30th June

Maps

- *map*: 2-cell embedding $G \rightarrow S$ of a graph G into a surface S .
- combinatorial description: $\mathcal{M} = (K; R, L)$, where
 - K : connected graph,
 - R : permutation of $D(K)$ (dart set of K),
 - L : involution of $D(K)$,
 - $\langle R, L \rangle$ transitive on $D(K)$.
- $\text{Aut}(\mathcal{M})$ acts semiregularly on $D(K)$.
- *regular map*: $\text{Aut}(\mathcal{M})$ acts transitively (i.e, regularly) on $D(K)$.

Cayley Maps

Let G be a group, Ω be a generating set s.t. $e_G \notin \Omega = \Omega^{-1}$ and p be a cyclic permutation of Ω .

- *Cayley graph* $C(G, \Omega)$:
 - vertex set: G ,
 - dart set: $G \times \Omega$
 - incidence: $g \xrightarrow{(g,x)} gx$
- *Cayley map* $CM(G, \Omega, p)$: $(C(G, \Omega); R, L)$, where
$$R(g, x) = (g, p(x)) \quad \text{and} \quad L(g, x) = (gx, x^{-1}).$$

t -balanced Cayley maps

Cayley map $\mathcal{M} = \text{CM}(G, \Omega, p)$ is

- *balanced* if $p(x^{-1}) = (p(x))^{-1}$ (1992 Širáň, Škoviera),
- *anti-balanced* if $p(x^{-1}) = (p^{-1}(x))^{-1}$ (1994 Škoviera, Širáň),
- *t-balanced* if $p(x^{-1}) = (p^t(x))^{-1}$ (1998 Martino, Schultz).

Necessarily, $t^2 \equiv 1 \pmod{|\Omega|}$.

Note: $\text{balanced} \Leftrightarrow 1\text{-balanced}$,
 $\text{antibalanced} \Leftrightarrow -1\text{-balanced}$.

Groups with sign structure

Širáň, Škovič 1992:

- *group with sign structure*: group G with a given subgroup G^+ of index at most 2,

Let G be a group with a generating set Ω .

- *sign structure induced by Ω* :

$$G^+ = \{g \in G; g = y_1 \dots y_{2n}, y_i \in \Omega, n \in \mathbb{N}_0\}$$

- *trivial sign structure*: $G = G^+$.

Theorem. Let $\mathcal{M} = CM(G, \Omega, p)$ be a regular t -balanced Cayley map. Then either Ω induces a nontrivial sign structure on G or \mathcal{M} is balanced.

Trivial sign structure

It can be shown, that canonical double cover of regular balanced Cayley map with trivial sign structure is regular balanced Cayley map with *nontrivial* sign structure.

Therefore, we can assume that sign structures induced by our maps are nontrivial. Equivalently, underlying graph is *bipartite*.

If $[G : G^+] = 2$ we say that G is a \mathbb{Z}_2 -*extension* of G^+ .

t -automorphisms

Martino, Schultz, Škovič :

Let $G = G^+(\theta, s)$ be a group with sign structure and $t \in \mathbb{Z} \setminus \{0\}$.

- t -automorphism: $\varphi \in \text{Sym}(G)$, $\varphi(x) \in G^+ \Leftrightarrow x \in G^+$ and

$$\varphi(xy) = \varphi(x)\varphi^{\pi(x)}(y),$$

$(\pi : G \rightarrow \{1, t\}, \pi(x) = 1, \text{ for } x \in G^+, \pi(x) = t, \text{ for } x \notin G^+)$.

Note: $t^2 \equiv 1 \pmod{\text{ord}(\varphi)}$.

Theorem. *Let $\mathcal{M} = CM(G, \Omega, p)$ be a Cayley map such that Ω induces a nontrivial sign structure on G . Then \mathcal{M} is regular and t -balanced if and only if there exists a t -automorphism φ of G whose restriction to Ω is equal to p .*

φ is called the rotary t -automorphism of \mathcal{M} .

Regular t -balanced Cayley maps

Regular t -balanced Cayley map \mathcal{M} is fully characterized by a triple (G, φ, x) where

- G is a group with sign structure,
- φ is a t -automorphism of G , and
- $x \in G$ is a member of a generating self-inverse orbit of φ .

New point of view

- underlying graph is bipartite,
- t is fixed, not necessarily $1 \leq t < |\Omega|$, i.e., 6-valent balanced map is also 7-balanced, -5 -balanced, etc.
- basic object is G^+ , instead of G .

We say that $\mathcal{M} = CM(G, \Omega, p)$ is **from** G and **over** G^+ .

G^+ as basic object: \mathbb{Z}_2 -extensions

Širáň, Škovič 1992:

Proposition. *Let G^+ be a group, $\theta \in \text{Aut}(G^+)$, $s \in G^+$ be such that*

$$\begin{aligned}\theta(s) &= s \\ \theta^2 &= \nu_s\end{aligned}$$

($\nu_s(g) = sgs^{-1}$). Then the set $G^+ \times \mathbb{Z}_2$ together with the binary operation \cdot defined by

$$(g, i)(h, j) = (g\theta^i(h)s^{ij}, i + j)$$

is a group denoted by $G^+(\theta, s)$. Moreover any \mathbb{Z}_2 -extension of G^+ is isomorphic to some $G^+(\theta, s)$.

G^+ as basic object: t -automorphisms

Líšková, M, Škoviera 2007:

Let $G = G^+(\theta, s)$ be a group with sign structure. Let φ be a t -automorphism of G , $\xi = \varphi|_{G^+}$ and $w = \varphi(e, 1)(e, 1)^{-1}$ ($e = e_{G^+}$). Then $\xi \in \text{Aut}(G^+)$, $w \in G^+$ and

$$\varphi(g, i) = (\xi(g)w^i, i).$$

We will write $\varphi \sim (\xi, w)$

Theorem. Let $G = G^+(\theta, s)$ be a group with sign structure. Let $\xi \in \text{Aut}(G^+)$, $w \in G^+$ be given, and put $\overline{w} = \xi^{-t}(\xi^{t-1}(w)\xi^{t-2}(w)\dots\xi(w)w)$. Then the mapping $\varphi : (h, i) \mapsto (\xi(h)w^i, i)$ is a t -automorphism of G iff:

$$\begin{aligned}\xi\theta(\overline{w}^{-1}s) &= ws \\ \xi\theta &= \nu_w\theta\xi^t = \nu_{ws}\theta^{-1}\xi^t.\end{aligned}$$

G^+ as basic object: regular t -balanced Cayley maps

Regular t -balanced Cayley map \mathcal{M} is fully characterized by a sextuple $(G^+, \theta, s, \xi, w, g)$ where $\theta, \xi \in \text{Aut}(G^+)$, $s, w, g \in G^+$ and

- $G = G^+(\theta, s)$ is a group with sign structure,
- mapping $\varphi \sim (\xi, w)$ is a t -automorphism of G , and
- $(g, 1)$ is a member of a generating self-inverse orbit of φ .

t -algebras

Universal algebra $\mathcal{A} = (G^+, \circ, e, {}^{-1}, \theta, s, \xi, w, g)$ of type $(2, 1, 0, 1, 0, 1, 0, 0)$ is a t -algebra if

- it satisfies all equations from the last three slides (they depend on t), and
- ξ is a permutation on G^+ ,

i.e, $(G^+, \circ, e, {}^{-1})$ is a group, $G = G^+(\theta, s)$ is a group with sign structure and $\varphi \sim (\xi, w)$ is a t -automorphism of G .

If $x = (g, 1)$ is a member of a generating self inverse orbit of φ , we say that \mathcal{A} *represents* the corresponding regular t -balanced Cayley map \mathcal{M} .

Product of regular t -balanced Cayley maps

Theorem. *Let $\mathcal{A} = \mathcal{A}_1 \times \mathcal{A}_2$ be a direct product of t -algebras. If \mathcal{A} represents a regular t -balanced Cayley map \mathcal{M} , then there exist regular t -balanced Cayley maps \mathcal{M}_1 and \mathcal{M}_2 such that \mathcal{A}_1 represents \mathcal{M}_1 and \mathcal{A}_2 represents \mathcal{M}_2 , respectively.*

We say that \mathcal{M} is a *product* of \mathcal{M}_1 and \mathcal{M}_2 and write

$$\mathcal{M} = \mathcal{M}_1 \odot \mathcal{M}_2$$

(this is a special case of the parallel product (Wilson 1994, Orbanic 2007)).

Decomposition results I

Proposition. *The direct product decompositions of a t -algebra $\mathcal{A} = (G^+, \circ, e, {}^{-1}, \theta, s, \xi, w, g)$ are in one to one correspondence with the inner product decompositions of the group G^+ into factors which are both θ - and ξ -invariant.*

Theorem. *Every regular t -balanced Cayley map*

from cyclic
 above a/an Abelian group
nilpotent

is a product of regular t -balanced Cayley maps

"from" cyclic
 above Abelian p -groups.
(nilpotent)

(If $2 \nmid |H|$, then "from" $H \Leftrightarrow$ from $H \times \mathbb{Z}_2$.)

Abelian groups

Lemma. *Let G^+ be an Abelian group of odd order (in the additive notation). Then every regular t -balanced Cayley map \mathcal{M} from $G = G^+ \times \mathbb{Z}_2$ can be represented by a t -algebra \mathcal{A} such that*

- 1) $\theta = \text{id}$ and $s = 0$,
- 2) $\xi^{t-1} = \text{id}$,
- 3) $(t+1) \times w = 0$ and
- 4) $\xi(w) = w$.

Lemma. *Let G^+ be a finite Abelian group and let $\psi \in \text{End}(G^+)$. Then there exists a positive integer n such that*

$$G^+ = \text{Ker } \psi^n \oplus \text{Im } \psi^n.$$

Abelian p -groups, p -odd

Lemma. *Let G^+ be an Abelian p -group of odd order, let $\mathcal{A} = (G^+, +, 0, -, \text{id}, 0, \xi, w, g)$ be a t -algebra representing regular t -balanced Cayley map \mathcal{M} and let $\psi = \xi - \text{id}$. Then, either \mathcal{M} is balanced or*

- 1) $G^+ = \text{Ker } \psi \oplus \text{Im } \psi$,
- 2) $\text{Ker } \psi = [w]$,
- 3) $\mathcal{M} = \mathcal{M}_a \odot \mathcal{M}_b$, where \mathcal{M}_a is over $[w]$ and \mathcal{M}_b is over $\text{Im } \psi$,
- 4) \mathcal{M}_a is antibalanced, and
- 5) \mathcal{M}_b is balanced.

Both \mathcal{M}_a and \mathcal{M}_b are t -balanced, too, i.e., the valency of \mathcal{M}_a divides $t + 1$ and the valency of \mathcal{M}_b divides $t - 1$.

Decomposition results II

Theorem. *Every regular t -balanced Cayley map \mathcal{M} from an Abelian group can be expressed as a product*

$$\mathcal{M} = \mathcal{M}_a \odot \mathcal{M}_b \odot \mathcal{M}_2,$$

where

- \mathcal{M}_a is antibalanced from $\mathbb{Z}_n \times \mathbb{Z}_2$,
- \mathcal{M}_b is balanced from $H \times \mathbb{Z}_2$, $|H|$ is odd, and
- \mathcal{M}_2 is from a 2-group.

(We allow some factors to be trivial, i.e., K_2 .)

Abelian 2-groups?

Proposition. *Any regular t -balanced Cayley map \mathcal{M} from \mathbb{Z}_{2^e+1} is antibalanced. Moreover, the valency of \mathcal{M} is one of 2^e ($e > 2$ two embeddings), 4 and 2.*

Theorem. *Every regular t -balanced Cayley map \mathcal{M} from \mathbb{Z}_{2^n} can be expressed as a product*

$$\mathcal{M} = \mathcal{M}_a \odot \mathcal{M}_b,$$

where \mathcal{M}_a is antibalanced from \mathbb{Z}_{2^k} , \mathcal{M}_b is balanced from \mathbb{Z}_{2^l} , $n = kl$ and $(k, l) \sim 1$.

Proposition. *Any regular t -balanced Cayley map \mathcal{M} from $\mathbb{Z}_{2^e} \times \mathbb{Z}_2$ with $G^+ = \mathbb{Z}_{2^e}$ has valency 2^e . Moreover it is antibalanced or $2^{e-1} - 1$ -balanced.*

Thank You